## **AMENDMENTS TO THE CLAIMS:**

Claims 21- 27 are canceled without prejudice or disclaimer. Claims 3-14, 19 are amended. The following is the status of the claims of the above-captioned application, as amended.

Claim 1.(Original) A method for deinking wastepaper comprising the steps of

i) pulping the wastepaper at a pH between 4 and 8.5 in the presence of deinking agent comprising a lipase and a fatty acid ester; and

ii) removing the thereby dislodged ink particles.

Claim 2.(Original) A method according to claim 1, wherein the consistency in step i) is from about 0.5% to about 15%.

Claim 3. (Currently amended.) A method according to any of the preceding claims claim 1, wherein the wastepaper comprises old newspapers (ONP).

Claim 4. (Currently amended.) A method according to claim 3, wherein the amount of ONP old newspapers constitutes at least 10% by weight of the total amount of wastepaper.

Claim 5. (Currently amended.) A method according to claim 4, wherein the wastepaper consists essentially of <u>old newspapers ONP</u>.

Claim 6. (Currently amended.) A method according to any of claims 1-2 claim 1, wherein the wastepaper comprises waste magazines (WM).

Claim 7. (Currently amended.) A method according to claim 6, wherein the amount of <del>WM</del> <u>waste</u> <u>magazines</u> constitutes at least 10% by weight of the total amount of wastepaper.

Claim 8. (Currently amended.) A method according to claim 7, wherein the wastepaper consists essentially of <del>WM</del> waste magazines.

Claim 9. (Currently amended.) A method according to any of claims 1-2 claim 1, wherein the wastepaper comprises ONP and WM old newspapers and waste magazines.

Claim 10. (Currently amended.) A method according to claim 9, wherein the wastepaper comprises 1-60% by weight of WM waste magazines and 40-99% by weight of ONP old newspapers.

Claim 12. (Currently amended.) A method according to any of the preceding claims claim 1, wherein the pulping with the deinking agent is carried out at a temperature from 25 to 75°C.

Claim 13. (Currently amended.) A method according to any of the preceding claims claim 1, wherein the fatty acid ester is a methyl ester, an ethyl ester, a n-propyl ester, an isopropyl ester, a n-butyl ester, an isobutyl ester, a sec-butyl ester, a tert-butyl ester, a monoglyceride, a diglyceride or a triglyceride of a  $C_6$ - $C_{22}$  fatty acid, the  $C_6$ - $C_{22}$  fatty acid being optionally substituted with one or more hydroxy, ethoxy, n-propoxy and/or isopropoxy groups.

Claim 14. (Currently amended.) A method according to any of claims 1-13 claim 1, wherein the fatty acid ester is a  $C_6$ - $C_{22}$  fatty acid, which has been alkoxylated with ethylene oxide, propylene oxide, or a combination thereof.

Claim 15. (Original) A method according to claims 13 or 14, wherein the fatty acid moiety of the fatty acid ester is selected from the group consisting of caproic acid (6:0), enanthic cid (7:0), caprylic acid (8:0) pelargonic acid (9:0), capric acid (10:0) undecylenic acid (11:0), lauric acid (12:0), tridecylic acid (13:0), myristic acid (14:0), palmitic acid (16:0), stearic acid (18:0), palmitoleic acid (16:1), oleic acid (18:1), elaidic acid (18:1), ricinoleic acid (18:1), linoleic acid (18:2), linolenic acid (18:3) and mixture thereof.

Claim 16. (Original) A method according to claims 13 or 14, wherein the fatty acid moiety of the fatty acid ester is substituted with one or more ethoxy and/or isopropoxy groups.

Claim 17. (Original) A method according to claim 16, wherein fatty acid moiety of the fatty acid ester is substituted with ethoxy and isopropoxy groups of the general formulae

-(O-CH<sub>2</sub>-CH<sub>2</sub>)<sub>x</sub>-(O-CH(CH<sub>3</sub>)-CH<sub>2</sub>)<sub>y</sub>-OH, -(O-CH<sub>2</sub>-CH<sub>2</sub>)<sub>x</sub>-(O-CH<sub>2</sub>-CH(CH<sub>3</sub>))<sub>y</sub>-OH, -(O-CH(CH<sub>3</sub>)-CH<sub>2</sub>)<sub>y</sub>-(O-CH<sub>2</sub>-CH<sub>2</sub>)<sub>x</sub>-OH, or  $-(O-CH_2-CH(CH_3))_y-(O-CH_2-CH_2)_x-OH,$ 

wherein x is an integer in the range from 1 to 25, and y is an integer in the range from 1 to 10.

Claim 18. (Original) A method according to claim 17, wherein x is an integer in the range from 1 to 10, and y is an integer in the range from 1 to 5.

Claim 19. (Currently amended.) A method according to any of claims 14-18 claim 14, wherein the fatty acid ester is a triglyceride.

Claim 20.( Original) A method according to claim 19, wherein the fatty acid ester is Hartaflot G-5000<sup>TM</sup>.

Claims 21-27 (Cancelled)